arXiv:1210.6217v1 [math.CO] 23 Oct 2012

REFLECTION GROUP RELATIONS ARISING FROM CLUSTER ALGEBRAS

AHMET I. SEVEN

Abstract. There is a well-known anology between cluster algebras and Kac-moody algebras: roughly speaking, Kac-Moody algebras are associated with symmetrizable generalized Cartan matrices while cluster algebras correspond to skew-symmetrizable matrices. In this paper, we study an interplay between these two classes of matrices. In particular, we obtain relations in the Weyl groups of Kac-Moody algebras that come from mutation classes of skew-symmetrizable matrices. These relations generalize those obtained by Barot and Marsh for finite type. As an application, we obtain some combinatorial properties of the mutation classes of skew-symmetrizable matrices.

1. Introduction

There is a well-known anology between cluster algebras and Kac-moody algebras: roughly speaking, Kac-Moody algebras are associated with symmetrizable generalized Cartan matrices while cluster algebras correspond to skew-symmetrizable matrices. In this paper, we study an interplay between these two classes of matrices. In particular, we obtain relations in the Weyl groups of Kac-Moody algebras that come from mutation classes of skew-symmetrizable matrices. These relations generalize those obtained by Barot and Marsh for finite type. As an application, we obtain some combinatorial properties of the mutation classes of skew-symmetrizable matrices.

To state our results, we need some terminology. Let us recall that an integer matrix B is skew-symmetrizable if DB is skew-symmetric for some diagonal matrix D with positive diagonal entries. For any matrix index k, the mutation of a skew-symmetrizable matrix B at k is another skew-symmetrizable matrix $\mu_k(B) = B'$:

$$B' = \begin{cases} B'_{i,j} = -B_{i,j} & \text{if } i = k \text{ or } j = k \\ B'_{i,j} = B_{i,j} + sgn(B_{i,k})[B_{i,k}B_{k,j}]_+ & \text{else} \end{cases}$$

(where we use the notation $[x]_+ = max\{x, 0\}$ and $sgn(x) = x/|x|$ with $sgn(0) = 0$). Mutation is an involutive operation, so repeated mutations give rise to the *mutation-equivalence* relation on skew-symmetrizable matrices. On the other hand, for any skew-symmetrizable $n \times n$ matrix B, a directed graph $\Gamma(B)$, called the diagram of B, is associated as follows: the vertices of $\Gamma(B)$ are the indices $1, 2, ..., n$ such that there is a directed edge from i to j if and only if $B_{ji} > 0$, and this edge is assigned the weight $|B_{ij}B_{ji}|$. Then the mutation μ_k can naturally be viewed as a transformation on diagrams (see Section 2 for a description). Note also that if

---

Date: October 23, 2012.

The author's research was supported in part by the Scientific and Technological Research Council of Turkey (TUBITAK).

B is skew-symmetric then the diagram $\Gamma(B)$ may be viewed as a quiver and the corresponding mutation operation is also called quiver mutation.

Given the appearance of the mutation operation in many different areas of mathematics, it is natural to study properties of the mutation classes of skew-symmetrizable matrices and the associated diagrams. In this paper, we consider mutation classes of skew-symmetrizable matrices with an acyclic diagram; here a diagram is called acyclic if it does not have any oriented cycles at all. To study such mutation classes, a useful notion has been the notion of a *quasi-Cartan companion.* More precisely a quasi-Cartan companion of B is a symmetrizable matrix A whose diagonal entries are equal to 2 and whose off-diagonal entries differ from the corresponding entries of B only by signs [1]. We will use a specific type of quasi-Cartan companions, which carry information about the the corresponding diagram. To be more explicit, let us recall that, for a skew-symmetrizable matrix B, by a subdiagram of $\Gamma(B)$ we always mean a diagram obtained from $\Gamma(B)$ by taking an induced (full) directed subgraph on a subset of vertices and keeping all its edge weights the same as in $\Gamma(B)$. By a cycle in $\Gamma(B)$ we mean a subdiagram whose vertices can be labeled by elements of $\mathbb{Z}/m\mathbb{Z}$ so that the edges betweeen them are precisely $\{i, i+1\}$ for $i \in \mathbb{Z}/m\mathbb{Z}$. We call a quasi-Cartan companion A of a skew-symetrizable matrix B *admissible* if it satisfies the following sign condition: for any cycle Z in $\Gamma(B)$, the product $\prod_{\{i,j\}\in Z}(-A_{i,j})$ over all edges of Z is negative if Z is oriented and positive if Z is non-oriented [6]. The main examples of admissible companions are the generalized Cartan matrices: if $\Gamma(B)$ is acyclic, i.e. has no oriented cycles at all, then the quasi-Cartan companion A with $A_{i,j} = -|B_{i,j}|$, for all $i \neq j$, is admissible, furthermore (for B with an acyclic diagram)

(1.1)

any matrix in its mutation class has an admissible quasi-Cartan companion.

This property has been shown in [7] for skew-symmetric matrices; for general skew-symmetrizable matrices it is conjectural.

These admissible companions can be considered as Gram matrices of a bilinear form with respect to particular bases. To be more explicit, let us also recall that, for a skew-symmetrizable B with an acylic diagram $\Gamma(B)$ and the corresponding generalized Cartan matrix A with $A_{i,i} = 2$ and $A_{i,j} = -|B_{i,j}|$ for $i \neq j$, there is an associated root system in the root lattice spanned by the simple roots α_i [5]. For each simple root α_i, the corresponding reflection $s_{\alpha_i} = s_i$ is the linear isomorphism defined on the basis of simple roots as $s_i(\alpha_j) = \alpha_j - A_{i,j}\alpha_i$. The group W generated by these (simple) reflections is called the Weyl group of the root system. Then the real roots are defined as the vectors obtained from the simple roots by the action of W, i.e. by a sequence of reflections. It is well known that the coordinates of a real root with respect to the basis of simple roots are either all nonnegative or all nonpositive, see [5] for details. For each real root $\alpha = w(\alpha_i)$, where $w \in W$, there is a corresponding reflection $s_\alpha = ws_iw^{-1}$. Following [2], for a skew-symmetrizable matrix B' which is mutation-equivalent to B with a quasi-Cartan companion A', we call a basis $\mathcal{B} = \{\beta_1, ..., \beta_n\}$ of the root lattice a companion basis if each β_i is a real root and $A' = (<\beta_i, \beta_j^\vee>)$, the matrix of the pairings between the roots and the coroots. We call an companion basis admissible if it can be obtained from the basis of simple roots by a sequence of mutations (Definition 2.1) and/or sign changes. Each admissible quasi-Cartan companion of B' has an admissible companion basis (Proposition 2.2).

For a skew-symmetrizable matrix B, we define, for vertices i, j of $\Gamma(B)$, the following:

$$m_{ij} = \begin{cases} 2 & \text{if } i \text{ and } j \text{ are not connected;} \\ 3 & \text{if } i \text{ and } j \text{ are connected by an edge of weight } 1; \\ 4 & \text{if } i \text{ and } j \text{ are connected by an edge of weight } 2; \\ 6 & \text{if } i \text{ and } j \text{ are connected by an edge of weight } 3. \\ \infty & \text{if } i \text{ and } j \text{ are connected by an edge of weight } \geq 4. \end{cases}$$

It is well-know that, if $\Gamma(B)$ is acyclic, then for the corresponding simple reflections s_i, the order of $s_i s_j$ in W is equal to m_{ij}. Our first main result gives the relations for the reflections of roots that form a companion basis for a matrix in the mutation class of B:

Theorem 1.1. *Suppose that B_0 is a skew-symmetrizable matrix whose diagram is acyclic. Suppose also that B is mutation-equivalent to B_0 and let A be an admissible quasi-Cartan companion of B. Let $\mathcal{B} = \{\beta_1, ..., \beta_n\}$ be a companion basis for A. Then, under the assumption* (1.1)*, the reflections $t_i := s_{\beta_i}$ satisfy the following relations:*

(i) *$t_i^2 = e$ for any vertex i*
(ii) *For any two vertices i, j in $\Gamma(B)$, $(t_i t_j)^{m_{ij}} = e$*
(iii) *For any oriented cycle $C = \{1, ..., d\}$ in Γ with*

$$1 \xrightarrow{w_1} 2 \xrightarrow{w_2} \cdots i-1 \xrightarrow{w_{i-1}} i \xrightarrow{w_i} i+1 \xrightarrow{w_{i+1}} \cdots \to d \xrightarrow{w_d} 1,$$

and any vertex i in C we have, writing $t_k = t_l$ if $k \equiv l \bmod d$,

$$(t_i t_{i+1} ... t_{i+d-1} t_{i+d-2} ... t_{i+1})^m = e$$

where, writing $q_j = \sqrt{w_j}$ and $x = (q_i ... q_{i+d-2} - q_{i+d-1})^2$ with $q_k = q_l$ if $k \equiv l \bmod d$, the number m is the following: if $x = 0$, then $m = 2$; if $x = 1$, then $m = 3$; if $x = 2$, then $m = 4$; if $x = 3$, then $m = 6$; if $x \geq 4$, then $m = \infty$.

If $\mathcal{B}$ is admissible, then $\{t_1, ..., t_n\}$ is a generating set for W. Also A has an admissible companion basis.

Let us note that these relations generalize those obtained in [2] for finite type. It has also been shown in [2] that, for finite type, these relations give a presentation of the corresponding finite Weyl group. We conjecture that the above relations provide a presentation of an infinite Weyl group W.

As an application of our study of the Weyl group in relation with the mutation operation, we obtain the following combinatorial property of the mutation classes.

Theorem 1.2. *Suppose that B is a skew-symmetrizable matrix such that $\Gamma(B)$ is acyclic. Suppose also that any edge in $\Gamma(B)$ has weight which is greater than or equal to 4. Then, under the assumption* (1.1)*, for any B' which is mutation-equivalent to B, any edge-weight in $\Gamma(B')$ is greater than or equal to 4.*

Our next result gives another combinatorial property of the mutation classes.

Theorem 1.3. *Suppose that B and C are skew-symmetrizable matrices whose diagrams $\Gamma(B)$ and $\Gamma(C)$ are acyclic and, in both diagrams, any edge has weight which is greater than or equal to 4. Suppose that, for both diagrams, the underlying undirected weightless graphs are equal. Let B' and C' be the matrices obtained from B*

and C by the same sequence of mutations respectively. Then, under the assumption (1.1)*, the underlying undirected weightless graphs of the diagrams $\Gamma(B')$ and $\Gamma(C')$ are equal.*

We prove our results in Section 3 after some preparation in Section 2.

2. Preliminaries

In this section, we will recall some more terminology and prove some statements that we will use to prove our results. First, let us recall that the diagram of a skew-symmetrizable (integer) matrix has the following property:

(2.1) the product of weights along any cycle is a perfect square, i.e. the square of an integer.

Thus we can use the term diagram to mean a directed graph, with no loops or two-cycles, such that the edges are weighted with positive integers satisfying (2.1). Let us note that if an edge in a diagram has weight equal to one, then we do not specify its weight in the picture.

For any vertex k in a diagram Γ, the associated mutation μ_k changes Γ as follows [3]:

- The orientations of all edges incident to k are reversed, their weights intact.
- For any vertices i and j which are connected in Γ via a two-edge oriented path going through k (see Figure 1), the direction of the edge $\{i, j\}$ in $\mu_k(\Gamma)$ and its weight γ' are uniquely determined by the rule

$$\pm\sqrt{\gamma} \pm \sqrt{\gamma'} = \sqrt{\alpha\beta}\,, \tag{2.2}$$

  where the sign before $\sqrt{\gamma}$ (resp., before $\sqrt{\gamma'}$) is "+" if i, j, k form an oriented cycle in Γ (resp., in $\mu_k(\Gamma)$), and is "−" otherwise. Here either γ or γ' can be equal to 0, which means that the corresponding edge is absent.
- The rest of the edges and their weights in Γ remain unchanged.



Figure 1. Diagram mutation

This operation is involutive, i.e. $\mu_k(\mu_k(\Gamma)) = \Gamma$, so it defines an equivalence relation on the set of all diagrams. More precisely, two diagrams are called *mutation-equivalent* if they can be obtained from each other by applying a sequence of mutations. The *mutation class* of a diagram Γ is the set of all diagrams which are mutation-equivalent to Γ. If B is a skew-symmetrizable matrix, then $\Gamma(\mu_k(B)) = \mu_k(\Gamma(B))$ (see Section 1 for the definition of $\mu_k(B)$). Let us note that if B is not skew-symmetric, then the diagram $\Gamma(B)$ does not determine B as there could be several different skew-symmetrizable matrices whose diagrams are equal; however, if a skew-symmetrizing matrix D is fixed, then $\Gamma(B)$ determines B, so mutation class of $\Gamma(B)$ determines that of B (the matrix $\mu_k(B)$ shares the same skew-symmetrizing matrix D with B [3]).

Let us recall that an $n \times n$ matrix A is called symmetrizable if there exists a diagonal matrix D with positive diagonal entries such that DA is symmetric. A crucial property of A is sign symmetry: $sgn(A_{i,j}) = sgn(A_{j,i})$. We say that A is a quasi-Cartan matrix if it is symmetrizable and all of its diagonal entries are equal to 2. A *quasi-Cartan companion* (or "companion" for short) of a skew-symmetrizable matrix B is a quasi-Cartan matrix A with $|A_{i,j}| = |B_{i,j}|$ for all $i \neq j$. The basic example of a quasi-Cartan companion of B is the associated generalized Cartan matrix A, which is defined as $A_{i,j} = -|B_{i,j}|$, for all $i \neq j$.

let us first discuss an extension of the mutation operation to quasi-Cartan companions:

Definition 2.1. Suppose that B is a skew-symmetrizable matrix and let A be a quasi-Cartan companion of B. Let k be an index. For each sign $\epsilon = \pm 1$, "the ϵ-mutation of A at k" is the quasi-Cartan matrix $\mu^{\epsilon}(A) = A'$ such that for any $i, j \neq k$: $A'_{i,k} = \epsilon sgn(B_{k,i})A_{i,k}$, $A'_{k,j} = \epsilon sgn(B_{k,j})A_{k,j}$, $A'_{i,j} = A_{i,j} - sgn(A_{i,k}A_{k,j})[B_{i,k}B_{k,j}]_+$. If A is admissible, then A' is a quasi-Cartan companion of $B' = \mu_k(B)$

Note that for $\epsilon = -1$, one obtains the formula in [1, Proposition 3.2]. Also note that if D is a skew-symetrizing matrix of B, then D is also a symmetrizing matrix for A, with $DA = S$ symmetric. If we consider S as the Gram matrix of a symmetric bilinear form on $\mathbb{Z}^n$ with respect to a basis $\mathcal{B} = \{e_1, ..., e_n\}$, then $DA' = S'$ is the Gram matrix of the same symmetric bilinear form with respect to the basis $\mathcal{B}' = \{e'_1, e'_2, ..., e'_n\}$ defined as follows: $e'_k = -e_k$; $e'_i = e_i - A_{k,i}e_k$ if $\epsilon B_{k,i} > 0$; $e'_i = e_i$ if else. We write $\mu_k(\mathcal{B}) = \mathcal{B}'$. If A is admissible and $\mathcal{B}$ is a companion basis, then $s_{e'_i} = s_{e_k}s_{e_i}s_{e_k}$, if $\epsilon B_{k,i} > 0$.

Proposition 2.2. *Suppose that B_0 is a skew-symmetrizable matrix whose diagram is acyclic and let A_0 be the corresponding generalized Cartan matrix. Suppose also that B is mutation-equivalent to B_0. Then any admissible quasi-Cartan companion A of B can be obtained from an admissible companion A'_0 of B_0 by a sequence of mutations. Furthermore A'_0 can be obtained from A, if necessary, by simultaneous sign changes in some rows and columns.*

These statements follow from the basic property that if A is an admissible companion of B, then $\mu^{\epsilon}(A) = A'$ is an admissible companion of $\mu_k(B)$, see [6] and [7].

3. Proofs of Main Results

First we will prove some lemmas that we use to prove our theorems.

Lemma 3.1. *In the set-up of Theorem 1.1, suppose that an edge $\{i, j\}$ in $\Gamma(B)$ has weight which is less than or equal to 4. Then the order of $s_{\beta_i}s_{\beta_j}$ is as stated in Theorem 1.1.*

Proof. Let us first note that for β_i in $\mathcal{B}$, the reflection s_{β_i} acts on the root lattice $V = span(\mathcal{B})$ as follows: $s_{\beta_i}(\beta_j) = \beta_j - A_{i,j}\beta_i$ for all $i, j = 1, ..., n$. Then the relations are obtained in the same way as for the groups generated by reflections in [4]. More explicitly, for an edge $\{i, j\}$ in $\Gamma(B)$ whose weight w is less than 4, the relations are obtained from a direct check; if $w \geq 4$, then $s_{\beta_i}s_{\beta_j}$ has infinite order on $span(\beta_i, \beta_j)$ as can be checked directly. □

Proof of Theorem 1.1: Let us first note that for β_i in $\mathcal{B} = \{\beta_1, ..., \beta_n\}$, the reflection s_{β_i} acts on the root lattice $V = span(\mathcal{B})$ as follows: $s_{\beta_i}(\beta_j) = \beta_j - A_{i,j}\beta_i$ for all $i, j = 1, ..., n$.

The relation (i) is well-known; (ii) follows from Lemma 3.1. As for the relation (iii), we assume, without loss of generality, that $i = 1$. Let $B' = \mu_2 ... \mu_{d-2}\mu_{d-1}(B)$ and, e.g., for $\epsilon = -1$, let $\mathcal{B}' = \mu_2^{\epsilon} ... \mu_{d-2}^{\epsilon}\mu_{d-1}^{\epsilon}(\mathcal{B})$. Then for β'_d we have $s_{\beta'_d} = s_{\beta_2} ... s_{\beta_{d-1}} s_{\beta_d} s_{\beta_{d-1}} ... s_{\beta_2}$ and $\beta'_1 = \beta_1$, furthermore the weight of the edge $\{1, d\}$ in $\Gamma(B')$ is equal to $(q_1 ... q_{d-1} - q_d)^2$. Thus, by Lemma 3.1, we have the relation (iii). The rest of the claims follow from Proposition 2.2.

Proof of Theorem 1.2: Let us suppose that B' is mutation-equivalent to B and $\Gamma(B)$ contains an edge $\{r, s\}$ whose weight is less than 4. Let t_r and t_s be the reflections corresponding to the companion basis elements (the companion basis can be obtained from the basis of simple roots by mutating accordingly). Then $3 \leq m_{r,s} \leq 6$, where $m_{r,s}$ is the order of $t_r t_s$ (Lemma 3.1).

To prove the theorem we will use a geometric representation of $\sigma : W \to GL(V)$ [4, Chapter 5.3]. More precisely, Let V be the n-dimensional real vector space with basis $\bar{\alpha}_1, \ldots \bar{\alpha}_n$ and equip V with the symmetric bilinear form $\bar{B}$ such that $\bar{B}(\bar{\alpha}_i, \bar{\alpha}_i) = 2$ and $\bar{B}(\bar{\alpha}_i, \bar{\alpha}_j) = -2\cos(\pi/m_{ij})$ for $i \neq j$. For each $\bar{\alpha}_i$, the corresponding reflection σ_i is defined on V as $\sigma_i : v \mapsto v - \bar{B}(v, \bar{\alpha}_i)\bar{\alpha}_i$; furthermore this action preserves the bilinear form $\bar{B}$. The elements of V that can be obtained from $\bar{\alpha}_1, \ldots \bar{\alpha}_n$ by a sequence of reflections are called the real roots. Then W, which is generated by $\{s_1, ..., s_n\}$, acts on V by $\sigma(s_i) = \sigma_i$. Furthermore, this action is faithful [4, Chapter 5.4], so the order of $\sigma(t_i)\sigma(t_j)$ is equal to the order of $t_i t_j$ for all i, j. Also $\sigma(t_i)$ is a reflection with respect to a real root $\bar{\beta}_i$, for all i. On the other hand, the assumptions on the weights of $\Gamma(B)$ implies that $\bar{B}(\bar{\alpha}_i, \bar{\alpha}_j) \neq 0$ if and only if $|\bar{B}(\bar{\alpha}_i, \bar{\alpha}_j)| = 2$; so $\bar{B}(\bar{\beta}_i, \bar{\beta}_j) = 2k$ for some integer k, implying that the order of $\sigma(t_r)\sigma(t_s)$ is 2 (if $k = 0$) or it is ∞ (if $k \neq 0$). This contradicts our assumption that $3 \leq m_{r,s} \leq 6$. This completes the proof.

Proof of Theorem 1.3: It is enough to prove the theorem assuming that $C_{i,j} = 2$ whenever $C_{i,j} \neq 0$. Let us suppose to the contrary that the conclusion of the theorem does not hold. Then we may assume, without loss of generality, that

(*) for $i = 1, ..., k-1$ and $B_i := \mu_i ... \mu_2\mu_1(B)$ and $C_i := \mu_i ... \mu_2\mu_1(C)$, the underlying graphs of $\Gamma(B_i)$ and $\Gamma(C_i)$ are equal,

however, for $B' = \mu_k(B_{k-1})$ and $C' = \mu_k(C_{k-1})$, the underlying graphs of $\Gamma(B')$ and $\Gamma(C')$ are not equal. Thus we may assume, without loss of generality, that

(**) $\Gamma(B')$ contains an edge $\{r, s\}$, however the vertices r and s are not connected in $\Gamma(C')$.

For any $\epsilon = \pm 1$, let $\mathcal{B}' = \mu_k^{\epsilon} ... \mu_2^{\epsilon}\mu_1^{\epsilon}(\mathcal{B})$, $\mathcal{C}' = \mu_k^{\epsilon} ... \mu_2^{\epsilon}\mu_1^{\epsilon}(\mathcal{C})$ be the companion bases obtained from the corresponding bases of simple roots $\mathcal{B} = \{\alpha_1, ..., \alpha_n\}$ and $\mathcal{C} = \{\bar{\alpha}_1, ..., \bar{\alpha}_n\}$. Let $\{t_1, ..., t_n\}$ and $\{\tau_1, ..., \tau_n\}$ be the reflections corresponding to the elements in $\mathcal{B}', \mathcal{C}'$ respectively. Then (*) implies that $\tau_i = \sigma(t_i)$, for all i, where σ is the (faithful) geometric representation that we discussed above. In particular, the order of $t_r t_s$ is equal to the order of $\tau_r \tau_s$. This contradicts our assumption (**) (which implies that the order of $t_r t_s$ is ∞, however the order of $\tau_r \tau_s$ is 2 by Lemma 3.1). This completes the proof.

References

[1] M. Barot, C. Geiss and A. Zelevinsky, Cluster algebras of finite type and positive symmetrizable matrices. J. London Math. Soc. (2) 73 (2006), no. 3, 545–564.

[2] M. Barot, R. Marsh, Reflection group presentations arising from cluster algebras, arXiv:1112.2300 (2011)

[3] S. Fomin and A. Zelevinsky, Cluster Algebras II, Inv. Math. 12 (2003), 335-380.

[4] J. Humphreys, Reflection groups and Coxeter groups, Cambridge University Press (1990).

[5] V. Kac, Infinite dimensional Lie algebras, Cambridge University Press (1991).

[6] A. Seven, Cluster algebras and semipositive symmetrizable matrices, Trans. Amer. Math. Soc. 363(5), 2011, 2733-2762.

[7] A. Seven, Cluster algebras and symmetric matrices, arXiv:1201.4270

[8] D. Speyer and H. Thomas, Acyclic cluster algebras revisited, arXiv:1203.0277, (2012).

Middle East Technical University, Department of Mathematics, 06800, Ankara, Turkey
E-mail address: `aseven@metu.edu.tr`